AAO Foundation Award Final Report

Secondary Investigators	Principal Investigator	
Secondary Investigators N/A Award Type Orthodontic Faculty Development Fellowship Award-AAO Restricte Gift Project Title Candidate Gene Studies of Mandibular Prognathism Using DNA Samples Acquired from Patients' Saliva Project Year 2008 Institution University of Pittsburgh Introduction: In orthodontics one of the most challenging aspects treating patients is the prediction of mandibular growth especia for patients that show more pronounced characteristics mandibular development. It is well documented that there is strong link of mandibular prognathism and genetics. Recent studi suggested a number of loci as linked to mandibular prognathis. The goal of this study was to test if any of these chromosom regions were associated with mandibular prognathism in a group subjects recruited at the University of Pittsburgh School of Den Medicine. Methods: The inclusion criteria for the Class III patient selection included cephalometric values indicative of Class III such as negatify the plane etc. Additional Class III criteria include negative OJ avisually prognathic profile as determined by the patient's clinic evaluation. The subjects were 44 prognathic/skeletal Class patients and 36 patients of orthognathic facial type serving as to control group. Single nucleotide polymorphism markers we selected in the chromosomal regions 1p22.1, 3q26.2, 5p13-p1 6q23, 11q22, 12q13.13, 12q23, and 19p13. Genotypes we generated using TaqMan chemistry. Chi-square, Fisher's exact tes were used to detect overrepresentation of alleles. Results: No association was found between the candidate region studied and mandibular prognathism. Conclusions: Although no associations between mandibul prognathism and genetic variation in candidate regions we detected, this project shows that larger scale projects including to collection of careful phenotypical clinical information and biologic samples is feasible in the orthodontic setting of the University	Co Investigator	Dr. Maria Tassopoulou-Fishell
Award Type Orthodontic Faculty Development Fellowship Award-AAO Restricte Gift Project Title Candidate Gene Studies of Mandibular Prognathism Using DNA Samples Acquired from Patients' Saliva Project Year 2008 Institution University of Pittsburgh Litroduction: In orthodontics one of the most challenging aspects treating patients is the prediction of mandibular growth especia for patients that show more pronounced characteristics mandibular development. It is well documented that there is strong link of mandibular prognathism and genetics. Recent studi suggested a number of loci as linked to mandibular prognathis. The goal of this study was to test if any of these chromoson regions were associated with mandibular prognathism in a group subjects recruited at the University of Pittsburgh School of Den Medicine. Methods: The inclusion criteria for the Class III patient selectic included cephalometric values indicative of Class III such as negati ANB, negative Witts appraisal (> -1.0 mm for men), positive A plane etc. Additional Class III criteria include negative OJ a visually prognathic profile as determined by the patient's clinic evaluation. The subjects were 44 prognathic/skeletal Class patients and 36 patients of orthognathic facial type serving as t control group. Single nucleotide polymorphism markers we selected in the chromosomal regions 1p22.1, 3q26.2, 5p13-p1 6q23, 11q22, 12q13.13, 12q23, and 19p13. Genotypes we generated using TaqMan chemistry. Chi-square, Fisher's exact tes were used to detect overrepresentation of alleles. Results: No association was found between the candidate region studied and mandibular prognathism. Conclusions: Although no associations between mandibul prognathism and genetic variation in candidate regions we detected, this project shows that larger scale projects including tollection of careful phenotypical clinical information and biologic samples is feasible in the orthodontic setting of the University		Dr. Alex Vieira
Award Type Orthodontic Faculty Development Fellowship Award-AAO Restricter Gift Project Title Candidate Gene Studies of Mandibular Prognathism Using DNA Samples Acquired from Patients' Saliva Project Year 2008 Institution University of Pittsburgh Introduction: In orthodontics one of the most challenging aspects treating patients is the prediction of mandibular growth especia for patients that show more pronounced characteristics mandibular development. It is well documented that there is strong link of mandibular prognathism and genetics. Recent studies suggested a number of loci as linked to mandibular prognathism. The goal of this study was to test if any of these chromoson regions were associated with mandibular prognathism in a group subjects recruited at the University of Pittsburgh School of Den Medicine. Methods: The inclusion criteria for the Class III such as negation and the plane etc. Additional Class III criteria include negative OJ at visually prognathic profile as determined by the patient's clinic evaluation. The subjects were 44 prognathic/skeletal Class patients and 36 patients of orthognathic facial type serving as to control group. Single nucleotide polymorphism markers were selected in the chromosomal regions 1p22.1, 3q26.2, 5p13-p1 6q23, 11q22, 12q13.13, 12q23, and 19p13. Genotypes were generated using TaqMan chemistry. Chi-square, Fisher's exact tes were used to detect overrepresentation of alleles. Results: No association was found between the candidate region studied and mandibular prognathism. Conclusions: Although no associations between mandibul prognathism and genetic variation in candidate regions we detected, this project shows that larger scale projects including to collection of careful phenotypical clinical information and biologic samples is feasible in the orthodontic setting of the University	Secondary Investigators	N/A
Project Title Candidate Gene Studies of Mandibular Prognathism Using DNA Samples Acquired from Patients' Saliva Project Year 2008 Institution University of Pittsburgh Summary/Abstract (250 word maximum) Introduction: In orthodontics one of the most challenging aspects treating patients is the prediction of mandibular growth especia for patients that show more pronounced characteristics mandibular development. It is well documented that there is strong link of mandibular prognathism and genetics. Recent studi suggested a number of loci as linked to mandibular prognathism. The goal of this study was to test if any of these chromosom regions were associated with mandibular prognathism in a group subjects recruited at the University of Pittsburgh School of Den Medicine. Methods: The inclusion criteria for the Class III patient selectic included cephalometric values indicative of Class III such as negatificational class III criteria include negative OJ at visually prognathic profile as determined by the patient's clinic evaluation. The subjects were 44 prognathic/skeletal Class patients and 36 patients of orthognathic facial type serving as to control group. Single nucleotide polymorphism markers we selected in the chromosomal regions 1p22.1, 3q26.2, 5p13-p16q23, 11q22, 12q13.13, 12q23, and 19p13. Genotypes we generated using TaqMan chemistry. Chi-square, Fisher's exact tes were used to detect overrepresentation of alleles. Results: No association was found between the candidate region studied and mandibular prognathism. Conclusions: Although no associations between mandibul prognathism and genetic variation in candidate regions we detected, this project shows that larger scale projects including t collection of careful phenotypical clinical information and biologic samples is feasible in the orthodontic setting of the University	Award Type	
Project Year 2008 Institution University of Pittsburgh Inroduction: In orthodontics one of the most challenging aspects for patients that show more pronounced characteristics mandibular development. It is well documented that there is strong link of mandibular prognathism and genetics. Recent studies suggested a number of loci as linked to mandibular prognathis. The goal of this study was to test if any of these chromoson regions were associated with mandibular prognathism in a group subjects recruited at the University of Pittsburgh School of Den Medicine. Methods: The inclusion criteria for the Class III patient selection included cephalometric values indicative of Class III such as negatificative of Class III such as negatificative of Class III patient selections and the plane etc. Additional Class III criteria include negative OJ as visually prognathic profile as determined by the patient's clinic evaluation. The subjects were 44 prognathic/skeletal Class patients and 36 patients of orthognathic facial type serving as to control group. Single nucleotide polymorphism markers we selected in the chromosomal regions 1p22.1, 3q26.2, 5p13-p1 6q23, 11q22, 12q13.13, 12q23, and 19p13. Genotypes we generated using TaqMan chemistry. Chi-square, Fisher's exact tes were used to detect overrepresentation of alleles. Results: No association was found between the candidate regions tudied and mandibular prognathism. Conclusions: Although no associations between mandibul prognathism and genetic variation in candidate regions we detected, this project shows that larger scale projects including to collection of careful phenotypical clinical information and biologic samples is feasible in the orthodontic setting of the University		Orthodontic Faculty Development Fellowship Award-AAO Restricted Gift
Project Year 2008 Institution University of Pittsburgh Introduction: In orthodontics one of the most challenging aspects treating patients is the prediction of mandibular growth especia for patients that show more pronounced characteristics mandibular development. It is well documented that there is strong link of mandibular prognathism and genetics. Recent studi suggested a number of loci as linked to mandibular prognathis. The goal of this study was to test if any of these chromosom regions were associated with mandibular prognathism in a group subjects recruited at the University of Pittsburgh School of Den Medicine. Methods: The inclusion criteria for the Class III patient selection included cephalometric values indicative of Class III such as negative ANB, negative Witts appraisal (> -1.0 mm for men), positive A plane etc. Additional Class III criteria include negative OJ at visually prognathic profile as determined by the patient's clinic evaluation. The subjects were 44 prognathic/skeletal Class patients and 36 patients of orthognathic facial type serving as t control group. Single nucleotide polymorphism markers we selected in the chromosomal regions 1p22.1, 3q26.2, 5p13-p1 6q23, 11q22, 12q13.13, 12q23, and 19p13. Genotypes we generated using TaqMan chemistry. Chi-square, Fisher's exact tes were used to detect overrepresentation of alleles. Results: No association was found between the candidate region studied and mandibular prognathism. Conclusions: Although no associations between mandibul prognathism and genetic variation in candidate regions we detected, this project shows that larger scale projects including t collection of careful phenotypical clinical information and biologic samples is feasible in the orthodontic setting of the University	Project Title	
Institution University of Pittsburgh Summary/Abstract (250 word maximum) Introduction: In orthodontics one of the most challenging aspects treating patients is the prediction of mandibular growth especia for patients that show more pronounced characteristics mandibular development. It is well documented that there is strong link of mandibular prognathism and genetics. Recent studi suggested a number of loci as linked to mandibular prognathis: The goal of this study was to test if any of these chromoson regions were associated with mandibular prognathism in a group subjects recruited at the University of Pittsburgh School of Den Medicine. Methods: The inclusion criteria for the Class III patient selecti included cephalometric values indicative of Class III such as negati ANB, negative Witts appraisal (> -1.0 mm for men), positive A plane etc. Additional Class III criteria include negative OJ at visually prognathic profile as determined by the patient's clinic evaluation. The subjects were 44 prognathic/skeletal Class patients and 36 patients of orthognathic facial type serving as t control group. Single nucleotide polymorphism markers we selected in the chromosomal regions 1p22.1, 3q26.2, 5p13-p1 6q23, 11q22, 12q13.13, 12q23, and 19p13. Genotypes we generated using TaqMan chemistry. Chi-square, Fisher's exact tes were used to detect overrepresentation of alleles. Results: No association was found between the candidate regios studied and mandibular prognathism. Conclusions: Although no associations between mandibul prognathism and genetic variation in candidate regions we detected, this project shows that larger scale projects including t collection of careful phenotypical clinical information and biologic samples is feasible in the orthodontic setting of the University		
Summary/Abstract (250 word maximum) Introduction: In orthodontics one of the most challenging aspects treating patients is the prediction of mandibular growth especia for patients that show more pronounced characteristics mandibular development. It is well documented that there is strong link of mandibular prognathism and genetics. Recent studi suggested a number of loci as linked to mandibular prognathis. The goal of this study was to test if any of these chromosom regions were associated with mandibular prognathism in a group subjects recruited at the University of Pittsburgh School of Dem Medicine. Methods: The inclusion criteria for the Class III patient selectic included cephalometric values indicative of Class III such as negative ANB, negative Witts appraisal (> -1.0 mm for men), positive A plane etc. Additional Class III criteria include negative OJ a visually prognathic profile as determined by the patient's clinic evaluation. The subjects were 44 prognathic/skeletal Class patients and 36 patients of orthognathic facial type serving as t control group. Single nucleotide polymorphism markers we selected in the chromosomal regions 1p22.1, 3q26.2, 5p13-p16q23, 11q22, 12q13.13, 12q23, and 19p13. Genotypes we generated using TaqMan chemistry. Chi-square, Fisher's exact tes were used to detect overrepresentation of alleles. Results: No association was found between the candidate region studied and mandibular prognathism. Conclusions: Although no associations between mandibul prognathism and genetic variation in candidate regions we detected, this project shows that larger scale projects including to collection of careful phenotypical clinical information and biologic samples is feasible in the orthodontic setting of the University	Project Year	2008
Summary/Abstract (250 word maximum) Introduction: In orthodontics one of the most challenging aspects treating patients is the prediction of mandibular growth especial for patients that show more pronounced characteristics mandibular development. It is well documented that there is strong link of mandibular prognathism and genetics. Recent studi suggested a number of loci as linked to mandibular prognathism. The goal of this study was to test if any of these chromosom regions were associated with mandibular prognathism in a group subjects recruited at the University of Pittsburgh School of Den Medicine. Methods: The inclusion criteria for the Class III patient selection included cephalometric values indicative of Class III such as negational ANB, negative Witts appraisal (> -1.0 mm for men), positive A plane etc. Additional Class III criteria include negative OJ avisually prognathic profile as determined by the patient's clinic evaluation. The subjects were 44 prognathic/skeletal Class patients and 36 patients of orthognathic facial type serving as the control group. Single nucleotide polymorphism markers we selected in the chromosomal regions 1p22.1, 3q26.2, 5p13-p16q23, 11q22, 12q13.13, 12q23, and 19p13. Genotypes we generated using TaqMan chemistry. Chi-square, Fisher's exact test were used to detect overrepresentation of alleles. Results: No association was found between the candidate region studied and mandibular prognathism. Conclusions: Although no associations between mandibul prognathism and genetic variation in candidate regions we detected, this project shows that larger scale projects including teleficial of called in the orthodontic setting of the University	Institution	
treating patients is the prediction of mandibular growth especia for patients that show more pronounced characteristics mandibular development. It is well documented that there is strong link of mandibular prognathism and genetics. Recent studi suggested a number of loci as linked to mandibular prognathis. The goal of this study was to test if any of these chromosom regions were associated with mandibular prognathism in a group subjects recruited at the University of Pittsburgh School of Den Medicine. Methods: The inclusion criteria for the Class III patient selection included cephalometric values indicative of Class III such as negatian ANB, negative Witts appraisal (> -1.0 mm for men), positive Aplane etc. Additional Class III criteria include negative OJ a visually prognathic profile as determined by the patient's clinic evaluation. The subjects were 44 prognathic/skeletal Class patients and 36 patients of orthognathic facial type serving as to control group. Single nucleotide polymorphism markers we selected in the chromosomal regions 1p22.1, 3q26.2, 5p13-p16q23, 11q22, 12q13.13, 12q23, and 19p13. Genotypes we generated using TaqMan chemistry. Chi-square, Fisher's exact tes were used to detect overrepresentation of alleles. Results: No association was found between the candidate region studied and mandibular prognathism. Conclusions: Although no associations between mandibul prognathism and genetic variation in candidate regions we detected, this project shows that larger scale projects including t collection of careful phenotypical clinical information and biologic samples is feasible in the orthodontic setting of the University		
	_	treating patients is the prediction of mandibular growth especially for patients that show more pronounced characteristics of mandibular development. It is well documented that there is a strong link of mandibular prognathism and genetics. Recent studies suggested a number of loci as linked to mandibular prognathism. The goal of this study was to test if any of these chromosomal regions were associated with mandibular prognathism in a group of subjects recruited at the University of Pittsburgh School of Dental Medicine. Methods: The inclusion criteria for the Class III patient selection included cephalometric values indicative of Class III such as negative ANB, negative Witts appraisal (> -1.0 mm for men), positive A-B plane etc. Additional Class III criteria include negative OJ and visually prognathic profile as determined by the patient's clinical evaluation. The subjects were 44 prognathic/skeletal Class III patients and 36 patients of orthognathic facial type serving as the control group. Single nucleotide polymorphism markers were selected in the chromosomal regions 1p22.1, 3q26.2, 5p13-p12, 6q23, 11q22, 12q13.13, 12q23, and 19p13. Genotypes were generated using TaqMan chemistry. Chi-square, Fisher's exact tests were used to detect overrepresentation of alleles. Results: No association was found between the candidate regions studied and mandibular prognathism. Conclusions: Although no associations between mandibular prognathism and genetic variation in candidate regions were detected, this project shows that larger scale projects including the collection of careful phenotypical clinical information and biological
Pittsburgh School of Dental Medicine. Were the original, This study was recently concluded and therefore the results have	Wone the onicinal	
	_	not been published. Publishing this study is definitely in our future
proposal realized? intentions. A manuscript will be prepared for submission to the American Journal of Orthodontics and Dentofacial Orthopedics.	-	intentions. A manuscript will be prepared for submission to the

Were the results published? If not, are there plans to publish? If not, why not?	
Have the results of this proposal been presented? If so, when and where? If not, are there plans to do so? If not, why not?	As I mentioned before, this is a recently concluded study and we have not yet published or presented anything related to our topic. We are planning to do so in the future in the AADR and AAO meetings.